

MAGNETIC AND MAGNETOCALORIC PROPERTIES OF $(Er_{1-x}Y_x)Co_2$ COMPOUNDS



István BALASZ-MURESAN, Izabela BALASZ, Liviu CHIONCEL* and Emil BURZO

Faculty of Physics, Babes-Bolyai University, RO-400084, Cluj-Napoca, Romania

*Institut für Physik, University Augsburg, 86135 Augsburg, Germany



To obtain information of physical properties of $(Er_{1-x}Y_x)Co_2$ compounds, were studied by X-rays, magnetic measurements and band structure calculations.

EXPERIMENTAL

INTRODUCTION

- The rare earth - transition metal intermetallic compounds R-TM (R - rare earth; TM - transition metal) exhibit a rich number of challenging physical phenomena [1]
- $ErCo_2$ is ferrimagnetically ordered compound with magnetic transition temperature $T_C \approx 35$ K [1]
- The yttrium will change the physical properties of the $ErCo_2$.

- $(Er_{1-x}Y_x)Co_2$ with $x \leq 0.3 \Rightarrow$ in an induction furnace, under high purity argon atmosphere.
- X-ray diffraction analyses \Rightarrow all the samples shows only one phase (Bruker D8 Advance AXS diffractometer with $Cu\ K\alpha$ radiation)
- Magnetic measurements \Rightarrow in magnetic fields $\mu_0 H \leq 12 \cdot 10^4$ Oe and $4.2 \leq T \leq 500$ K (Oxford Instruments)



RESULTS AND DISCUSSION



- XRD $\Rightarrow (Er_{1-x}Y_x)Co_2$ compounds with $x \leq 0.3$ crystallize in a cubic $MgCu_2$ -type structure [1]. The lattice parameters, determined at room temperature, show a Veguad-type dependence.




Fig. 1. Cubic $MgCu_2$ -type structure of $ErCo_2$




Fig. 2. Lattice parameters of $Er_{1-x}Y_xCo_2$

- Magnetic measurements (1) \Rightarrow The temperature dependences of magnetizations of zero field cooled (ZFC) and field cooled (FC) $Er_{0.7}Y_{0.3}Co_2$ compound is given in Fig.3. The compounds are ferrimagnetically order - Fig.4.




Fig. 3. Temperature dependences of magnetizations in a field of 0.5 T, for sample $Er_{0.9}Y_{0.1}Co_2$ field cooled (FC) and zero field cooled (ZFC).




Fig. 4. Magnetization isotherms of $Er_{0.9}Y_{0.1}Co_2$

- Magnetic measurements (2) \Rightarrow The cobalt moments determined from saturation data ($H \leq 12$ T, $T = 4.2$ K) decrease when Y content is higher- Fig. 5. The reciprocal susceptibilities follows non-linear characteristic for a ferrimagnetic system - Fig. 6.




Fig. 5. Co moments in $Er_{1-x}Y_xCo_2$




Fig. 6. Reciprocal susceptibilities in $Er_{1-x}Y_xCo_2$

- Magnetic measurements (3) \Rightarrow Assuming that the effective erbium moment is given by its free ion value [2], the contributions of cobalt atoms to the Curie constants were determined. The effective cobalt moments, $M_{eff}(Co)$, decrease little in the investigated composition range - Fig.7. The ratio $r = S_p/S_0$ between the number of spins determined from effective cobalt moment and saturation one can be founded in spin fluctuation ($r \propto T_C^{-2/3}$) model - Fig 8.




Fig. 7. Composition dependences of the effective cobalt moments and Curie temperatures




Fig. 8. The dependence of the ratios $r = S_p/S_0$ on the Curie temperatures

- Magnetic measurements (4) \Rightarrow The computed entropy changes, ΔS , for the $Er_{0.1}Y_{0.9}Co_2$ compound is plotted in - Fig. 9. The $-\Delta S_{max}$ values follow a $H^{2/3}$ type [3] dependence as expected in mean field model - Fig. 10. The specific renormalized power in a field $H < 2.25$ T an only constant for a given composition decrease from 60 J/kg*T ($x=0$) to 12 J/kg*T ($x=0.2$).




Fig. 9. Magnetocaloric effect for $Er_{0.9}Y_{0.1}Co_2$




Fig. 10. $-\Delta S_{max}$ for $Er_{1-x}Y_xCo_2$



CONCLUSION

The substitution of erbium with yttrium in $ErCo_2$ decrease the magnetic interaction and also the magnetocaloric effect!!!



REFERENCES

- [1] E. Burzo, A. Chelkovski & H. R. Kirchmayr, Compounds of rare-earth elements and 3d elements. Landolt Börnstein Handbook, vol.19d2, Springer Verlag, (1990).
- [2] E. Burzo, Int. J. Magn. 3, 171 (1972)
- [3] Q.Y. Dong, H.W. Zhang, J.L. Shen, J.R. Sun, B.G. Shen, J. Magn. Magn. Mater. 319, 56 (2007)

Acknowledgements

This work was supported by the Romanian Ministry of Education and Research (UEFISCDI), grant no. PN- II- ID- PCE-2012- 4- 0028.